

RESEARCH ARTICLE

Distribution of Virulence Genes in *Salmonella* Serovars isolated from poultry farms in Kwara State, Nigeria

Ahmed O. Akeem^{1,*}, P. H Mamman², M. A. Raji¹, C. N. Kwanashie², I. A. Raufu¹ and A. Aremu³

¹Department of Veterinary Microbiology, University of Ilorin, Ilorin, Kwara State, Nigeria

²Department of Veterinary Microbiology, Ahmadu Bello University, Zaria, Kaduna State, Nigeria

³Department of Veterinary Pharmacology and Toxicology, University of Ilorin, Ilorin, Kwara State, Nigeria

Received: 01/06/2017; Accepted: 04/10/2017

Abstract: This study was aimed at investigating the prevalence and detection of virulence genes in *Salmonella* isolated from poultry farms in Kwara State, Nigeria. Ten commercial layer farms located in three senatorial districts of the state were sampled. The presence of *Salmonella* was investigated using standard bacteriological techniques, the isolates obtained were serotyped and polymerase chain reaction (PCR) was used to detect presence of two genes (*fimA* and *sefC*) in the isolates. Of the 300 samples collected, 24 (8.0%) were positive for *Salmonella*; 4 different serovars were identified; *Salmonella* Larochelle, *S. Muenster*, *S. enterica* ser. 6, 7:d, *S. enterica* ser. 45:d:1,7 and *Salmonella* Typhimurium, the most frequent being *Salmonella* Typhimurium (3%). Varying distribution patterns of the studied genes were observed among the isolates. While, *fimA* was found in all the 24 (100%) isolates, *sefC* was found only among the *Salmonella enterica* ser. 45: d: 1,7. The study indicates that the *fimA* gene is widely distributed among *Salmonella* irrespective of the serovars. However, the *sefC* gene appears to be serovar specific. The prevalence of *Salmonella* in chicken and poultry farms constitutes a major public health concern. Hence, further epidemiological studies are necessary.

Keywords: *Salmonella*, Kwara, Virulence, Poultry farms.

INTRODUCTION

Poultry industry is one of the major components of the Nigerian economy, serving as a source of income for farmers and a main source of high quality protein for the ever-growing population in Nigeria, due to their affordability and acceptability (Fagbamila *et al.*, 2010; Bettridge *et al.*, 2014). Poultry production plays an important role in the provision of animal protein, this account for about 25% of local meat production in Nigeria (Agbaje *et al.*, 2010). The

poultry industry in Nigeria has been expanding rapidly in past years despite facing many challenges like the global financial crisis, diseases outbreak such as Avian influenza, inadequate credit facilities, low egg production, poor chick quality, poor and low performing breeds, poor weight gain/ feed conversion, feeding and management issues (FAO, 2008; Agada *et al.*, 2014a). The Nigerian poultry industry improved from 150,700 million chickens in 2005 to 192,313 million in 2010 (FAO, 2015). Across different regions of the country, the poultry sector is characterized by a low level of production and weak levels of specialization (FAO, 2008).

Salmonellosis is an important health problem and a major challenge affecting both humans and animals worldwide. *Salmonella* species are recognized as the most prevalent agent of food poisoning. Salmonellosis in human is often associated with gastroenteritis, which is usually self-limiting. In some cases, particularly in children, pregnant women, infants, the elderly, and immunocompromised patients, *Salmonella* infection can lead to invasive and focal infections that can be severe (Hald *et al.*, 2007). The sources of nontyphoidal *Salmonella* are numerous, but most incidences of human foodborne illnesses have been traced to the consumption of poultry meat, eggs, and poultry-based food products (Rabsch *et al.*, 2001; Andrews and Bäumlner, 2005). The fact that these pathogens infect the host animals without apparent sickness contributes to high numbers of human infections (Patrick *et al.*, 2004; Dawoud *et al.*, 2011). Modes of transmission and carrier sources of nontyphoidal *Salmonella* are poorly understood in Africa due to the lack of

*Corresponding Author's Email: ahmed.oa@unilorin.edu.ng

coordinated national epidemiological surveillance systems (Kagambèga *et al.*, 2013).

In poultry, salmonellosis has become an important problem worldwide, causing great economic losses through substantial morbidity, mortality and drop in egg production in addition to huge amounts spent on vaccination and medication (Muhammed *et al.*, 2010; Agada *et al.*, 2014b). Poultry salmonellosis is related to host adapted serovars and remains a major constraint on poultry production in all parts of Nigeria (Garba *et al.*, 2013 and Mamman *et al.*, 2014). Salmonellosis in chicks is usually due to *Salmonella Pullorum*, the consequence of which is high mortality of both broilers and commercial layers (Idowu *et al.*, 2017). It is characterized by nonspecific signs such as weakness, depression, dehydration, loss of appetite, drooping wings, huddling, somnolence and ruffled feathers. Labored breathing or gasping, as well as diarrhea and pasting of the vent feathers may also be seen (Nógrády *et al.*, 2003). Most signs of salmonellosis in adult birds are thought to be due to *Salmonella Gallinarum* (Mamman *et al.*, 2014). In addition, zoonotic serovars of *Salmonella* are harbored in the gastrointestinal tracts of poultry with no apparent signs of illness. Hence, these *Salmonella* can be excreted in faeces by apparently healthy animals and may be transferred to raw foods of animal origin through contamination during slaughtering and processing (Sanchez *et al.*, 2002).

The virulence of *Salmonella* species is associated with a combination of chromosomal and plasmid factors (Oliveira *et al.*, 2003). The pathogenicity of an organism in a host is a function of its virulence. Some genes participate in adhesion and invasion such as *pef* (plasmid encoded fimbriae), *spv* (*Salmonella* plasmid virulence) *inv* (invasion gene) or *fim* (fimbriae gene) (Alaa and Adnan, 2011). Others play pivotal roles in the survival of organism in the host system- *mgtC* (Magnesium transport C) (Blanc-Potard and Groisman 1997) or in the actual manifestation of pathogenic processes for example *stn* (*Salmonella* toxin), *pip* A, B, D (Marcus *et al.*, 2000). Bacterial adherence (Kurkkonen *et al.*, 1993) is considered to be a prerequisite for infection, and there are evidences that many bacteria have surface processes, such as fimbriae (Clegg *et al.*, 1985; Aslanzadeh and Paulissen, 1992) or pili, that facilitate attachment to particular receptors on the epithelial cell

surface, to urinary tract or intestinal mucus. Although certain strains of *E. coli* fimbriae are known to be important for virulence, only type 1 fimbriae (*fimA*) have been implicated in *Salmonella* pathogenicity.

Control of *Salmonella* infection will not only culminate into expansion of poultry industry in Nigeria but also lead to reduction in the threat of foodborne illness due to *Salmonella* in human. Effective prevention and control measures cannot be undertaken unless the status of the disease and epidemiology are well investigated. Therefore, the present study was undertaken to investigate the prevalence and detect the virulence genes in *Salmonella* serovars in selected layer farms in Kwara state.

MATERIALS AND METHODS

Study area

The study was carried out in Kwara State, Nigeria. Kwara State is located between latitudes (8° 30'N) and longitudes (5° 00'E). The state shares a common internal boundary with Niger State in the North, Kogi State in the East, Oyo, Ekiti and Osun States in the South and an international boundary with the Republic of Benin in the west. Kwara State is a summer rainfall area, with an annual rainfall range of 1,000 mm to 1,500 mm. The rainy season begins at the end of March and lasts until early September, while the dry season begins in early October and ends in early March. Temperature is uniformly high and ranges between 25 °C and 30 °C in the wet season throughout the season except in July – August when the clouding of the sky prevents direct insolation (heatstroke) while in the dry season it ranges between 33 °C to 34 °C (NBS, 2010).

Study strategy

The study involved sampling for *Salmonella* in layers farms that agreed to participate in the study. Two Local Governments from each of the three senatorial districts of the state were selected based on their commercial poultry production according to data obtained from the state veterinary services. The Local Governments were Ilorin South, Ilorin-West, Moro, Offa, Irepodun and Asa. Twelve farms were selected randomly by balloting for the study, two farms per Local Government; however, two farms declined.

Sample collection

Layer farms were visited across the selected Local Governments, and the following samples were collected: cloacal swabs, litters composed of faeces mixed with saw dust (from deep litters system), non-medicated water and poultry feeds. Twenty-seven (27) cloacal swabs were taken per farm with one each from a bird, litters (n=1/farm), water (n=1/farm) and feed (n=1/farm) making a total of 300 samples. Cloacal swabs were collected using sterile swabs, 25 g of wet litters, approximately 50 g of feed and 100 ml of untreated water were collected per farm. All the samples were labelled properly and kept in a cool box containing ice packs and transported, within five (5) hours, to Veterinary microbiology laboratory, University of Ilorin for analysis.

Sample Processing

Samples were pre-enriched in buffered peptone water (Lab M, Lancashire, UK) at the ratio of 1:10 sample to broth incubated at 37 °C for 24 hours. 1 ml of the pre-enriched broth was inoculated in 9 ml of Selenite-F (Oxoid, Hampshire, UK) and Rappaport-Vassiliadis (Oxoid, Hampshire, UK) incubated for 24 hours at 37 °C and 42 °C, respectively (ISO, 2002; OIE, 2012).

Isolation, Identification, and Serotyping of *Salmonella*

The selective broths (Selenite F and Rappaport-Vassiliadis) were subcultured onto *Salmonella*-

Shigella agar (Oxoid, Hampshire, UK) and Xylose Lysine Deoxycholate agar (Oxoid, Hampshire, UK) and were incubated aerobically at 37 °C for 24 hours. The presumptive *Salmonella* isolates on XLD (pink/red with/without central black spot due to hydrogen sulphide production) and SSA (transparent colonies with black centre) were then subjected to standard biochemical tests (Andrews, 1992; MacFaddin, 2002). Isolates biochemically identified as *Salmonella* were streaked on nutrient agar slants and shipped to the WHO National *Salmonella* and *Shigella* Center, Bangkok, Thailand, for serotyping according to the Kauffmann-White Scheme (Raufu *et al.*, 2013; Issenhuth-Jeanjean *et al.*, 2014).

Polymerase Chain Reaction (PCR) analysis of the isolates

DNA extraction

DNA extraction was carried out in accordance with manufacturer's instruction using ZR fungal/bacterial DNA miniPrep™ (Zymo Research Corp. USA) at Department of Veterinary public health laboratory, University of Ilorin.

Primers

The sets of primer pairs used in the PCR assay are as shown in table 1. The primers were synthesized at Inqaba biotech (QN 201502628) South Africa.

Table 1: Primers sequence for PCR amplification.

Primer	Sequence: F-(5'-3') R-(3'-5')	Product size (bp)	Reference
<i>fim A</i>	F CCT TTC TCC ATC GTC CTG AA R CA CGA TCC GTC TAT TGT TGG	670	Huguet <i>et al.</i> , 1996
<i>Sef C</i>	F GCGAAAACC AAT GCG ACT GTA R CCCACC AGA AAC ATT CAT CCC	1103	Murugkar <i>et al.</i> , 2003

PCR assay

PCR analysis was carried out at Bioscience Laboratory unit of International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria. The PCR reaction contained a final volume of 25 μ L which included 12.5 μ l master mix (Inqaba Biotech, S. Africa) containing (5 U/ μ l) Taq DNA polymerase, 2.5 mM each of dATP, dCTP, dTTP and dGTP, 50mM MgCl₂ and PCR buffer, 1.25 μ l (1 μ M) each of forward and reverse primers and 2.5 μ l of template DNA (100ng/ μ l DNA) and 7.5 μ l of deionized water. The cocktail was used for each sample. The reaction was performed in a thermocycler (Perkin-Elmer, USA) in 40 cycles of initial denaturation (94 °C for 5 minutes), denaturation (94 °C for 30 seconds), primer annealing (55 °C for 45 seconds) and primer extension (72 °C for 45 seconds) followed by final extension at 72 °C for 7 minutes. Fifteen microliters (15 μ l) of each PCR products were loaded on 2.0% agarose gel containing 0.5 μ l/ml ethidium bromide (Pharmacia, Sweden). The gel picture was captured using Gel doc 2000 documentation system (Pharmacia, Sweden).

RESULTS

Isolation of *Salmonella*

Of the 300 samples collected, 24 (8.0 %) were positive for *Salmonella*. The different *Salmonella* serovars and their rates of isolation from different sources are as shown in table 2. *Salmonella* Typhimurium was the most frequently isolated serovar from 2 (20 %) of feed and water samples each and 5(1.9 %) of the cloacal swabs. *Salmonella* Muenster was only isolated from 2 water samples (20 %).

Detection of *Salmonella* genes using PCR

In this study, PCR assay was performed for the detection of two virulence genes. The *fimA* gene was detected in all the isolates by demonstrating the presence of 670 bp PCR product (Figure 1.0). The *sefC* gene was detected by the presence of 1103 bp PCR product (Figure 2.0) only in 20.8 % (5/24) of the isolates confirmed to be *Salmonella enterica* ser. 45:d:1,7 serovars isolates (Table 3).

Table 2: Different *Salmonella* serovars and rates of their isolation from different sources.

Serovars	Sample Source				
	Cloacal swabs	Litters	Feeds	Water	Total
Larochele	0(0)	1(10.0)	0(0)	1(10.0)	2(0.7)
Muenster	0(0)	0(0)	0(0)	2(20.0)	2(0.7)
<i>S. enterica</i> ser. 6,7 d	3(1.1)	0(0)	3(30.0)	0(0)	6(2.0)
<i>S. enterica</i> ser. 45,d 1,7	2(0.7)	1(10.0)	2(20.0)	0(0)	5(1.7)
Typhimurium	5(1.9)	0(0)	2(20.0)	2(20.0)	9(3.0)
Total	10(3.7)	2(20.0)	7(70.0)	5(50.0)	24(8.0)
No. of samples examined	270	10	10	10	300

Table 3: Distribution of *fim A* and *sef C* genes among *Salmonella* isolated from poultry farms.

Serotype	No. of isolates	No. positive for virulence genes	
		<i>fim A</i>	<i>sef C</i>
<i>S. Larochele</i>	2.0	2.0	0
<i>S. Muenster</i>	2.0	2.0	0
<i>S. enterica</i> ser. 6,7 d	6.0	6.0	0
<i>S. enterica</i> ser. 45,d 1,7	5.0	5.0	5.0
<i>S. Typhimurium</i>	9.0	9.0	0
Total	24.0	24.0	5.0





Figure 1: Detection of *fimA* gene by PCR. Lane M-(50 bp) DNA ladder molecular weight marker, lane N-negative control (deionized water), 2 to 24-Test isolates.




Figure 2: Detection of *sefC* gene PCR. Lane M-(50 bp) DNA ladder molecular weight marker, lane N-negative control (deionized water), 1 to 24-Test isolates.

DISCUSSION

Salmonellosis is one of the major zoonotic foodborne diseases worldwide (Kagambèga *et al.*, 2013) and its prevalence in animals poses a continuous threat to man (Muhammad *et al.*, 2010). Several *Salmonella* serovars were isolated from all the sample sources examined in this study and many of these *Salmonella* serovars are known to be pathogenic to man. The isolation rate of *Salmonella* serovars in this study correlates with earlier study by Muhammad *et al.* (2010) which reported isolation rate of 9.0 % from farms in Jos, Plateau state, Nigeria. It is however higher than 2.8 % reported in Caribbean countries (Adesiyun *et al.*, 2014)

Salmonella Typhimurium was the most frequent isolated serovar and this agree with previous work which also reported this serovar as one of the common serovars isolated from poultry in Nigeria (Orji *et al.*, 2005). The isolation of invasive *Salmonella* such as serotype Typhimurium and other pathogenic salmonellae in this study highlights the risk of health hazards to consumers as contaminated poultry and poultry products may serve as source of infection to human food chain (Doaa, 2013).

Polymerase chain reaction (PCR) has become a potentially powerful alternative technique in microbiological diagnostics due to the fact that it is fast, simple, accurate, and reproducible (Pickup *et al.*, 2003). In this present study, *fimA* gene was detected in all the 24 *Salmonella* isolates. These finding is in agreement with earlier reports by Huguette *et al.* (1996), Alaa and Adnan (2011) which both found that the *fimA* gene contains sequences unique to *Salmonella* isolates and demonstrated that this gene is suitable for PCR target for the detection of *Salmonella* strains. Bacterial adherence (Kurkkonen *et al.*, 1993) is considered to be a baseline requirement for infection, and there is evidence that many bacteria have surface appendages, such as pili or fimbriae (Clegg *et al.*, 1985; Aslanzadeh and Paulissen, 1992), that facilitate their binding to specific receptors on the epithelial cell surface, or intestinal mucus. Although certain types of *Escherichia coli* fimbriae are known to be important for virulence, only type 1 fimbriae (*fimA*) have been implicated in *Salmonella* pathogenicity.

Salmonella Enteritidis fimbriae 14 (SEF14) is encoded by the *sef* operon, which

contains *sefC* gene. It contains 4 major protein subunits SefA, SefB, SefC, and SefD. SEF14 plays important role in the ability of *Salmonella* to colonize Peyer's patches and in the adhesion and invasion of epithelial cells of the host intestine (Castilla *et al.*, 2006). In the present study, the *sefC* gene was detected in 5(20.8 %) of the isolates which were confirmed to be *Salmonella enterica* ser. 45:d:1,7 by serotyping. The present finding does not agree with the findings of Rahman (1999) and Murugkar *et al.* (2003) which showed that except for strains of *S. enteritidis* and *S. gallinarum*, none of the other serotypes (*S. Typhimurium*, *S. Newport*, *S. Kentucky*, *S. Weltevreden* and *S. Indiana*) tested were found to contain *sefC* gene. These differences might be due to the fact that the serotype shown to contain this gene in our present study was not part of the serovars studied by Rahman (1999) and Murugkar *et al.* (2003) or might be due to evolution of new gene in this serovar. Future studies are needed to confirm this hypothesis.

CONCLUSION

Contamination of chicken farms with *Salmonella* indicate generally poor infrastructure and low biosecurity measures in farms in the state. The study reaffirmed the applicability of PCR amplification of *fimA* for identification of *Salmonella* species and reported for the first time the presence of *SefC* gene in *Salmonella* ser. 45:d:1,7.

ACKNOWLEDGEMENT

We would like to thank all laboratory technologists at the Department of Veterinary Microbiology, University of Ilorin and Bioscience Unit, IITA, Ibadan for the technical assistance.

Conflicts of interest: none

REFERENCES

- Adesiyun, A., Webb, L., Musai, L., Louison, B., Joseph, G., StewartJohnson, A., Samlal, S. and Rodrigo, S. (2014). Survey of *Salmonella* contamination in chicken layer farms in three Caribbean countries. *Journal of Food Protection*, **77**: 1471-1480.
- Agada, G. O. A., Abdullahi, I. O., Aminu, M., Odugbo, M., Chollom, S. C., Kumbish, P. R. and Okwori, A. E. J. (2014a). Prevalence and Antibiotic Resistance Profile of *Salmonella* Isolates from Commercial Poultry and Poultry

- Farm-handlers in Jos, Plateau State, Nigeria. *British Microbiology Research Journal*, **4**: 462-479.
- Agada, G. A., Abdullahi, I. O., Aminu, M., Odugbo, M., Chollom, S. C., Okeke, L. A. and Okwori, A. E. J. (2014b); Prevalence and risk factors associated with *Salmonella* species contamination of commercial poultry farms in Jos, Plateau State, Nigeria. *World Journal of Biology and Biological Science*, **2**: 49-61.
- Agbaje, M., Davies, R., Oyekunle, M. A., Ojo, O. E., Fasina, F. O. and Akinduti P. A. (2010) Observation on the occurrence and transmission pattern of *Salmonella* Gallinarum in commercial poultry farms in Ogun State, South Western Nigeria. *African Journal Microbiological Research*, **4**: 796-800.
- Alaa, A. J. and Adnan, H. A. (2011). Detection of *fimA* and *fimC* genes of *Salmonella* isolates by using Polymerase Chain Reaction. *Journal of Basrah Researches (Sciences)*, **37**: 27-36.
- Andrews, H. L. and Bäumler, A. J. (2005). *Salmonella* species. Pages 327–339 in Foodborne Pathogens: Microbiology and Molecular Biology. P. M. Fratamico, A. K. Bhunia, and J. L. Smith, ed. Caister Academic Press, Norwich, UK.
- Andrews, W. (1992). *Manuals of food quality control*. 4. Microbiological analysis. FAO food and nutrition paper 14/4, *Food and Agriculture Organization, Rome* **1**: 27-48.
- Aslanzadeh, J. and Paulissen, L. J. (1992). Role of the type 1 and type 3 fimbriae adherence and pathogenic of *Salmonella enteritidis* in mice. *Microbiology and Immunology*, **36**: 351-359.
- Bettridge, J.M., Lynch, S.E., Brena, M.C., Melese, K., Dessie, T., Terfa, Z.G., Desta, T.T., Rushton, S., Hanotte, O., Kaiser, P., Wigley, P. and Christley, R.M. (2014). Infection-interactions in Ethiopian village chickens. *Preventive Veterinary Medicine*, **117** (2): 358-366. DOI: 10.1016/j.prevetmed.2014.07.002
- Blanc-Potard, A. B. and Groisman, E. A. (1997) The *Salmonella selC* locus contains a pathogenicity island mediating intramacrophage survival. *EMBO Journal*, **16**: 5376-85.
- Castilla, K. S., Astolfi, C. S., Moreno, A. M., Nunes, I. A., and Piantino, F. A. J. (2006). Distribution of virulence genes *sefC*, *pefA* and *spvC* in *Salmonella* Enteritidis phage type 4 strains isolated in Brazil. *Brazilian Journal of Microbiology*, **37**: 135-139.
- Clegg, S., Pruckler, J. and Purcell, B. K. (1985). Complementation analyses of recombinant plasmids encoding type 1 fimbriae of members of the family *Enterobacteriaceae*. *Infection and Immunology*, **50**: 338–340.
- Dawoud, T. M., Herrera, P., Hanning, I., Kwon, Y. M. and Rieke, S. C. (2011). In vitro invasion of laying hen ovarian follicles by *Salmonella* Enteritidis strains. *Poultry Science*, **90**: 1134-1137.
- Doaa, M. A. E. (2013). Detection of *Salmonella typhimurium* in retail chicken meat and chicken. *Asian Pacific Journal of Tropical Biomedicine*, **3**(9): 678-681.
- Fagbamila, I., Kabir, J., Abdu, P., Omeiza, G., Ankeli P., Ngulukun, S., Maryam, M. and Jarlath, U. (2010). Antimicrobial screening of commercial eggs and determination of Tetracycline residue using two microbiological methods. *International Journal of Poultry Science*, **9**: 959-962.
- Food and Agricultural Organization (2008). Assessment of the Nigerian poultry market chain to improve biosecurity. Available: <http://www.fao.org/3/a-ak778e.pdf>.
- Food and Agricultural Organization (FAO) (2015). Data Source: FAOSTAT | FAO Statistics Division 2015 | December 2015.
- Garba A., Turaki, U. A., Abdurrahman, A., Goji, J. N., Bolajoko, I.J. Barde, Ahmed A., Sa'Adatu, I. and Abdullahi, A. S. (2013). The Threat of Salmonellosis to Commercial Poultry Production in Adamawa State, Nigeria. *Sokoto Journal of Veterinary Sciences*, **8**: 50-52.
- Hald, T., Lo Fo Wong, D. M. and Aarestrup, F. M. (2007). The attribution of human infections with antimicrobial resistant *Salmonella* bacteria in Denmark to sources of animal origin. *Food borne Pathogen and Disease*, **4**: 313-326.
- Huguet, J., Cohen, Subbaiah, M., Mechanda and Wei, L. (1996). PCR amplification of the *fimA* gene sequence of *Salmonella* Typhimurium, a specific method for detection of *Salmonella* spp. *Applied and Environmental microbiology*, **62**: 4303-4308.
- Idowu O. F., Lisa B., Marzia M., Jacob K., Sati S. N., Paola Z., Antonia A. L., Monica L., Paul, A. A., Junaidu, K., Jarlath, U., Antonia, R. and Maryam, M. (2017). *Salmonella* serovars and their distribution in Nigerian commercial chicken layer farms. *PLoS ONE*, **12**(3): e0173097.doi:10.1371/journal.
- International Organization of Standardization (ISO) 6579 (2002). Microbiology general guidelines on methods for the detection of *Salmonella*. International organization of standardization, Geneva, Switzerland
- Issenhuth-Jeanjean, S., Roggentin, P., Mikoleit, M., Guibourdenche, M., de Pinna, E., Nair, S., Fields, I. P. and Weill, F. (2014) . Supplement 2008-2010 (no. 48) to the White-Kauffmann-Le Minor scheme. *Research Microbiology*, **165**: 526-530.
- Kagambèga, A., Lienemann, T., Aulu, L., Traoré, A. S., Barro, N., and Siitonen, A. (2013). Prevalence and characterization of *Salmonella enterica* from the feces of cattle, poultry, swine and hedgehogs in Burkina Faso and their comparison to human *Salmonella* isolates. *BMC Microbiology*, **13**: 253-56.

- Kurkkonen, M. T., Raunio, T., Virkola, R., Lahtemaki, K., Makela, P. H., Klemm, P., Clegg, S. and Korhonen T. K. (1993). Basement membrane carbohydrate as a target for adhesion binding of type 1 fimbriae of *Salmonella* Enteritidis and *Achillea* to lamini. *Molecular Microbiology*, **7**: 227-237.
- MacFaddin, J. F. (2002). Biochemical tests for identification of medical bacteria. Williams and Wilkins, New York
- Mamman, P. H., Kazeem, H. M., Raji, M. A., Nok, A. J. and Kwaga, J. K. (2014). Isolation and Characterization of *Salmonella* Gallinarum from Outbreaks of Fowl Typhoid in Kaduna State, Nigeria. *International Journal of Public Health and Epidemiology*, **3**: 082-088
- Marcus, S. L., Brummell, J. H., Pfeifer, C. G. and Finlay, B. B. (2000). *Salmonella* pathogenicity islands: big virulence in small packages. *Microbial Infection*, **2**: 145-156
- Muhammad, M., Muhammad, L. U., Ambali, A., Mani, A. U., Azard, S., and Barco, L. (2010). Prevalence of *Salmonella* associated with chick mortality at hatching and their susceptibility to antimicrobial agents. *Veterinary Microbiology*, **140**: 131-135.
- Murugkar, H.V., Rahman, H. and Dutta, P. K. (2003). Distribution of virulence genes in *Salmonella* serovars isolated from man & animals. *Indian Journal of Medicine Research*, **117**: 66-70.
- National Bureau of Statistics (NBS) (2010). Commercial Agriculture Development Project (CADP),. NBS/CADP Baseline Survey Report. **5**: 5-46.
- Nógrády, N., Imre, A., Rychlik, I., Barrow, P.A. and Nagy, B., (2003). Growth and colonisation suppression of *Salmonella enterica* serovar Hadar in vitro and in vivo. *FEMS Microbiology Letters*, **218**: 127-133.
- Office International des Epizooties (OIE) (2012). Fowl typhoid and pullorum disease. In: Terrestrial manual. Office International des Epizooties, Paris, France, 3-5.
- Oliveira S. D., Rodenbusch C. R., Michael G. B., Cardoso M. I. R., Canal C. W. and Brandelli A. (2003). Detection of virulence genes in *Salmonella* Enteritidis isolates from different sources. *Brazilian Journal of Microbiology*, **34**: 123-124.
- Orji, M. U., Onuigbo, H. C., and Mbata, T. I. (2005). Isolation of *Salmonella* from poultry droppings and other environmental sources in Awka , Nigeria. *International Journal of Infectious Diseases*, **9**: 86–89.
- Patrick, M. E., Adcock, P. M., Gomez, T. M., Altekruze, S. E., Holland, B. H., Tauxe, R. V. and Swendlow, D. L. (2004). *Salmonella* Enteritidis infections, United States, 1985–1999. *Emerging Infectious Diseases*, **10**: 1-7.
- Pickup, R. W., Rhodes, G. and Hermon-Taylor, J. (2003). Monitoring bacterial pathogens in the environment: advantages of a multilayered approach. *Current Opinion Biotechnology*, **14**: 319-325.
- Rabsch, W., Tschäpe, H. and Bäumler, A. J. (2001). Non-typhoidal salmonellosis: Emerging problems. *Microbes and Infection*, **3**: 237-247.
- Rahman H. (1999). Prevalence of enterotoxin gene (*stn*) among different serovars of *Salmonella*. *Indian Journal of Medicine Research*, **110**: 43-6.
- Raufu, I., Bortolaia, V., Svendsen, C. A., Ameh, J. A., Ambali, A. G., Aarestrup, F. M. and Hendriksen, R. S. (2013). The first attempt of an active integrated laboratory-based *Salmonella* surveillance programme in the north-eastern region of Nigeria. *Journal of Applied Microbiology*, **115**: 1059-1067.
- Sanchez, S., Hofacre, C. L., Lee, M. D., Maurer, J. J. and Doyle, M. P. (2002). Animal sources of salmonellosis in humans. *Journal of American Veterinary Association*, **211**: 492-497.