

Medicinal Plants in Management of Diabetes Mellitus: An Overview

D.S.N.K. Liyanagamage¹, S. Jayasinghe^{1,*}, A.P. Attanayake² and V. Karunaratne¹

¹Department of Chemistry, Faculty of Science, University of Peradeniya, Peradeniya, Sri Lanka.

²Department of Biochemistry, Faculty of Medicine, University of Ruhuna, Sri Lanka.

Received: 20/07/2019 ; Accepted: 15/01/2020

Abstract: Diabetes mellitus has become a global epidemic over the past few decades and is one among the major causes of death worldwide. Although various modern medicines are in the market, herbal medicines have usually maintained worldwide popularity in achieving the primary health care needs of the diabetic population because of the multiple adverse effects related to them. Herbal medicines have a prolonged history of being used as medications and sources of medication for diabetes mellitus. Some of them are enclosed within the modern therapeutic arsenal of medicine, and others are used as complementary therapy or as dietary supplements by patients with diabetes. The present review describes on traditional antidiabetic herbal medicines with an emphasis on scientific investigations conducted on antidiabetic medicinal plants, polyherbal mixtures, isolated compounds and their associated toxicities.

Keywords: Antidiabetic compounds, antidiabetic medicinal plants, diabetes mellitus, polyherbal mixtures, toxicity of antidiabetic agents.

INTRODUCTION

Diabetes mellitus has become a devastating health problems nowadays and millions of people are fighting with this dilemma worldwide. Diabetes mellitus is an endocrine disease characterized by hyperglycaemia which occurs as a result of the inability of the pancreas to secrete insulin, defects in insulin action or both (ADA, 2013). Hyperglycaemia results in both microvascular and macrovascular complications which lead to long-term failure of various organs (Chawla *et al.*, 2016). Although numerous synthetic antidiabetic drugs are available, these medications have limitations in their use due to unwanted side effects (Chaudhury *et al.*, 2017). Hence, herbal products have been accepting a great demand for the management of diabetes due to low cost, high availability and less side effects.

The dependence of human beings on medicinal plants dates back to the beginning of life. Medicinal plants include various types of plants used for numerous medicinal purposes (Jamshidi-Kia *et al.*, 2018). During the past two decades, medicinal plants have become a topic of global importance. World health organization (WHO) estimated that 80% of world population relies on medicinal plants (Gopl *et al.*, 2014). In spite of the presence of

effective modern allopathic medicines, herbal medicines have often maintained global popularity in meeting the primary health care needs. Concurrently, many people in developed countries have begun to turn to alternative or complementary therapies with medicinal plants. However, there is a paucity of scientific research data on medicinal plants.

According to WHO reports, around 21,000 plant species have the potential for being used as medicinal plants (Modak *et al.*, 2007). There have been significant developments in the pharmacological evaluation of various medicinal plants used in traditional systems of medicine in the treatment of many of the chronic diseases. Diabetes mellitus is one of them. Importantly, the treatments using medicinal plants are considered safe with little or no side effects independent of the age or gender of the patients. Furthermore, medicinal plants are considered as rich resources of pharmacophores which can be used in drug development. In addition, medicinal plants play critical roles as important sources of nutrition.

The present review is an overview of diabetes mellitus, traditional medicines used in the management of diabetes mellitus, summary of antidiabetic medicinal plants, medicinal plants grown in Sri Lanka with scientifically proven antidiabetic activity, polyherbal mixtures with proven antidiabetic activity and toxicities of antidiabetic medicinal plants and polyherbal mixtures.

Diabetes mellitus has become a global epidemic and is considered as one of the emerging global health problems. The prevalence of diabetes has rapidly increased over the past few decades (WHO, 2016). Patients with diabetes have a high risk of morbidity and mortality than the healthy population. Globally, according to the International Diabetic Federation (IDF) Atlas, there were 425 million people with diabetes worldwide in 2016 (Ogurtsova *et al.*, 2017).

Many efforts have been devoted to search and develop optimal therapeutic agents for the management of diabetes mellitus. Due to various side effects associated with currently available oral antidiabetic drugs and their high cost, most of the diabetic patients especially in developing countries tend to use herbal plant remedies as alternative therapies for the management of diabetic mellitus (Ekor, 2014).

*Corresponding Author's Email: susanthij@pdn.ac.lk

 <https://orcid.org/0000-0002-7964-2596>



Traditional herbal medicines for the management of diabetes mellitus

The use of traditional medicine continues to play a crucial role in the healthcare delivery systems of many parts of the world. Indeed, according to WHO, 40 % of the total population relies on traditional medical systems directly or indirectly (Mazid *et al.*, 2012), while 80% of the population in developing countries depend almost exclusively on traditional medical practices, more specifically, plant medicines for their primary health care needs (Karunamoorthi *et al.*, 2013). Reports from developed countries also indicate that several diabetic patients are gradually turning to herbal remedies as alternatives to modern synthetic drugs (Rani *et al.*, 2015). The literature on medicinal plants with hypoglycaemic activity is vast. More than 1200 species of plants have been used ethno-pharmacologically to treat for diabetes mellitus (Delaviz *et al.*, 2017). Traditional systems of medicines always have played a significant role as therapeutic aid for the management of diabetes mellitus. In South East Asia, there is a vast volume of literature on traditional medicine. Generally, civilization has adopted broad cultural aspects which led to create different types of traditional medicine systems for over 3000 years. There are four main systems of traditional medicine in Sri Lanka namely; Ayurveda, Siddha, Unani and Deshiya Chikitsa (Waisundara and Watawana, 2014). The Ayurveda and Deshiya Chikitsa

systems use mainly plants and herbal preparations for the treatment of diabetes mellitus as single plants or plant mixtures. Sri Lankan traditional physicians regularly use many antidiabetic plants such as *Momordica charantia* (Bitter melon), *Allium sativum* (garlic), *Annona squamosa* (sugar apple), *Gymnema sylvestre* (Gymnea), *Syzygium cumini* (Malabar Plum), *Allium cepa* (onion) *etc.* for their daily prescriptions.

Natural products derived from medicinal plants play an important role as useful tools in pharmacological studies in the management of diabetes mellitus (Farzaei *et al.*, 2015). Researchers have proved that the antidiabetic activity of medicinal plants is due to the presence of phenolic compounds, flavonoids, terpenoids, coumarins and other ingredients which demonstrate hypoglycaemic activity (Aswathy and Jessykutty, 2017). In many cases, the detailed mechanisms of action of traditionally used antidiabetic plants are unknown or inconclusive (Patel *et al.*, 2012). As such, many studies are being conducted to elucidate the mechanisms of action of different plants and natural compounds (Grover *et al.*, 2002). Medicinal plant extracts are mainly responsible for regenerating the function of pancreatic tissues by elevating the insulin output or inhibiting the intestinal absorption of glucose or by facilitating metabolites in insulin dependent processes (Patel *et al.*, 2012). Some widely used antidiabetic medicinal plants that were recently scrutinized by scientific investigations are summarized in Table 1.

Table 1: A summary of antidiabetic plants which were recently scrutinized by scientific investigations.

Family	Plant	plant part	Extract	Type of Study	Reference
Alangiaceae	<i>Alangium lamarckii</i>	Leaf	Methanol	Streptozotocin nicotinamide induced rats	Kumar <i>et al.</i> , 2011
Alliaceae	<i>Allium sativum</i>	Bulb	Ethanol	Normal and streptozotocin-induced rats	Eidi <i>et al.</i> , 2006
	<i>Annona squamosa</i>	Leaf	Aqueous	Streptozotocin-nicotinamide induced rats	Shirwaikar <i>et al.</i> , 2004
Apocynaceae	<i>Catharanthus roseus</i>	Leaf	Aqueous	Streptozotocin-induced diabetic rats	Rasineni <i>et al.</i> , 2010
	<i>Tabernanthe iboga</i>	Root, bark	Aqueous	Rat pancreatic islets	Souza <i>et al.</i> , 2011
Caesalpinaceae	<i>Bauhinia variegata</i>	Stem bark	Aqueous	Alloxan induced hyperglycaemic rats	Kumar <i>et al.</i> , 2012
Cornaceae	<i>Cornus officinalis</i>	Fruit	Aqueous	Streptozotocin Induced diabetic rats	Han <i>et al.</i> , 2014
Cucurbitaceae	<i>Coccinia grandis</i>	Leaf	Aqueous	Streptozotocin-diabetic rats	Attanayake <i>et al.</i> , 2016
Fabaceae	<i>Medicago sativa</i>	Seed	Aqueous	Alloxan induced diabetic rats.	Helal <i>et al.</i> , 2013
Gentianaceae	<i>Enicostemma littorale</i>	Plant	Aqueous	Alloxan induced diabetic rats	Maroo <i>et al.</i> , 2003
Ginkgoaceae	<i>Ginkgo biloba</i>	Plant	Aqueous	Streptozotocin-induced diabetic rats	Cheng & Li 2012
Liliaceae	<i>Aloe vera</i>	Plant	Aqueous	Streptozotocin induced diabetic rats	Bahmani <i>et al.</i> , 2014
	<i>Asparagus racemosus</i>	Root	Ethanol	Streptozotocin-induced diabetic rats	Hannan <i>et al.</i> , 2012
Leguminosae	<i>Bougainvillea spectabilis</i>	Bark	Ethanol	Alloxan- induced diabetic rats	Jawla <i>et al.</i> , 2012
Rutaceae	<i>Aegle marmelos</i>	Seed	Aqueous	Diabetic rat	Kesari <i>et al.</i> , 2006
	<i>Syzygium cumini</i>	Seed	Ethyl acetate	Streptozotocin- induced diabetic rats	Kumar <i>et al.</i> , 2013

Sri Lankan medicinal plants with antidiabetic activity

The majority of rural population in Sri Lanka use indigenous medicinal systems for their primary health care. Considerable numbers of patients with diabetes mellitus have frequently used herbal supplements in the management of diabetes mellitus (Ediriweera and Ratnasooriya, 2009). However, only a few scientific studies have been reported on detailed antidiabetic mechanisms of medicinal plants in Sri Lanka. The studies on *P. embelica*, *P. marsupium*, *C. auriculata* confirmed the antihyperglycaemic effects of the selected plant extracts via the retardation of carbohydrate metabolizing enzymes (Abesundara et al., 2004). The decoctions of Sri Lankan medicinal plants as *Salacia reticulata* (Celastraceae) *Aegle marmelos* (Rutaceae) and *Momordica charantia* (Cucurbitaceae) have shown potent hypoglycaemic activity in laboratory animals (Karunanayake et al. 1984). Fernando et al (1991) studied the hypoglycaemic activity of some Sri Lankan plants including *Artocarpus heterophyllus*, *Bambusa vulgaris* and *Osbeckia octandra*. The above plant extracts exerted potent antidiabetic activity *in vivo*. The stem bark of *Gmelina arborea*, *Kokoona zeylanica*, *Spondias pinnata*, *Syzygium caryophyllatum*, arial parts of *Scoparia dulcis*, *Sida alnifolia*, and roots of *Languas galanga* and leaves of *Coccinia grandis* were also evaluated on antihyperglycaemic activity by Attanayake et al., (2013a, 2016b). Furthermore, Arambewela et al., (2005) scrutinized the antidiabetic activities of the aqueous and ethanolic extracts of *Piper betle* leaves in Wistar rats. A report by Arambewela et al., (2004) revealed the hypoglycaemic and antihyperglycaemic activities of the aqueous and the ethanolic extracts of *Alpinia calcarata* rhizomes in rats. Moreover, Sudasinghe et al., (2018) reported on hypoglycaemic and hypolipidaemic activities of the aqueous leaf extract of *Passiflora suberosa*. In addition to the work on the above Sri Lankan higher medicinal plants, antidiabetic compounds namely, zeorin, methyl β -orcinolcarboxylate and methylorsellinate have been isolated from the Sri Lankan lichen *Cladonia* species (Karunaratne et al., 2014; Thadhani and Karunaratne, 2017)

Antidiabetic compounds reported from medicinal plants

Bioactive compounds isolated from medicinal plants are in much demand than the synthetic compounds as antidiabetic drug leads owing to rich availability, efficacy and less side effects. Among the drugs approved in the therapy of diabetes in the last ten years, 49% are derived from plants (Gothai et al., 2016). More than 1200 antidiabetic compounds from medicinal plants have been discovered by scientists all over the world (Trojan-Rodrigues et al., 2012). The antidiabetic activities of isolated compounds could be varied based on mechanisms of their actions for lowering glucose levels including the inhibition of glucose absorption, improvement in insulin resistance and pancreatic functions (Chang et al., 2013)

Quercetin isolated from the aqueous extract of *Matricaria recutita* has shown high inhibitory effects

against glucosidase as yeast α -glucosidase, rat intestinal sucrase, and amylase with IC_{50} values as 8.86 μ g/mL, 216 μ mol/L, and 71 μ mol/L respectively (Yin et al., 2014). Mahanine obtained from the fruit pulp of *Murraya koenigii* showed α -glucosidase inhibitory activity as 21.4 \pm 0.4 μ mol/L compared with acarbose (IC_{50} = 15.2 \pm 0.6 μ mol/L) (Uvarani et al., 2014). Furthermore, mangiferin (IC_{50} = 299.7 \pm 42.3 μ mol/L) was isolated from 70% aqueous ethanolic leaf extracts of *Aquilaria sinensis* and showed significant α -glucosidase inhibitory activity than acarbose (IC_{50} = 576.2 \pm 58.5 μ mol/L) (Feng et al., 2011).

The major steps in the discovery of bioactive compounds from plants are solvent extraction, bioassays, isolation, characterization, toxicological evaluation, preclinical and clinical investigation of isolated bioactive compounds (Sasidharan et al., 2011). Table 2 summarizes some common antidiabetic compounds that have been isolated from medicinal plants.

Polyherbal formulations for the management of diabetes mellitus

Medicinal plants with ethnomedicinal value are currently being screened for their therapeutic potential. Herb-herb combinations are basically found in polyherbal therapy and have been used in Ayurvedic medicine for thousands years (Che et al., 2013). In the Ayurvedic literature 'Sarangdhar Samhita', the concept of polyherbalism and combined extracts of plants are chosen rather than individual ones for the treatment of diabetes mellitus in the traditional system of medicine (Parasuraman et al., 2014). The Unani medicine has also achieved global acceptance due to the amazing clinical efficiency of the polyherbal formulations (Shailajan, 2015). In the majority of traditional systems, diabetes has been successfully managed by the polyherbal mixtures instead of a single herb because of the powerful synergism and less side effects (Yuan et al., 2017). Combinations often lead to promising effects in the management of diseases over a single drug. Naturally occurring herbs and herbal ingredients are organized into certain formulas that offer potential interactions such as mutual enhancement, mutual assistance, mutual restraint and mutual antagonism (Alabi et al., 2017). Many scientific studies have discovered that polyherbal mixtures produce a greater effect, as compared to individual plant extracts. Combined plant extracts also focus on multiple targets at the same time to provide greater effectiveness (Briskin, 2000). In the preparation of polyherbal mixtures, the extent of individual plants used is less in achieving desirable pharmacological action than in a single herbal preparation. Hence, it reduces the risk of harmful side-effects. In addition, polyherbal formulations also prevent the need to take more than one different single herbal formulation at a time. Therefore, all these benefits has led to the popularity of polyherbal mixtures among patients with diabetes. Many researchers have studied polyherbal mixtures in order to evaluate their effectiveness in acute and long term interventions. These studies have revealed that combined plant mixtures have a wide therapeutic range due to the presence of different phytochemicals. Most of the polyherbal mixtures have shown effectiveness even at low or high doses without any toxic effects (Parasuraman

et al., 2014). A summary of recent studies on polyherbal formulation used in the management of diabetes mellitus is shown in Table 3.

Table 2: Reported antidiabetic active compounds isolated from medicinal plants.

Plant	Used part	Active compound	Extract	Mode of study	References
Acacia pennata	Shoot tips	Polyphenols, caffeic acid	Aqueous	In vitro	Wongsa et al., 2012
Bauhinia forficata	Leaves, flower	Kaempferol-3-neohesperidoside	Ethanollic	Rat soleus muscle	Brahmachari, 2011
Capparis moonii	Fruits	Gallotannins	Ethanollic	L 6 cells	Kanaujia et al., 2010
Curcuma longa	Rhizome	Curcumin, demethoxycurcumin, bisdemethoxycurcumin, ar-turmerone	Ethanollic	Diabetic mice and in vitro	Kuroda et al., 2005
Eleutherine americana	Bulb	Eleutherinoside A	Methanol Ethanol	Compound isolation	Ieyama et al., 2011
Macaran gatanarius	Seeds	Ellagitannins (mallotinic acid, corilagin, chebulagic acid, macatannins A and B)	Ethanol	Compound isolation	Gunawan-Puteri and Kawabata, 2010
Murraya koenigii	Leaves	Mahanibine	Petroleum ether	Diabetic rats	Dinesh kumar et al., 2010
Piper retrofractum	Fruits	Piperidine alkaloids: piperine, pipernonaline, dehydropipernonaline	Ethanollic	3T3-L1 adipocytes	Kim et al., 2011
Salacia oblonga	Root, stem, leaves	Salacinol, kotalanol, mangiferin	Aqueous	Compound isolation,	Muraoka et al., 2011
Swertia punicea	Whole plant	Methylswertianin, bellidifolin	Ethanollic	Diabetic mice	Tian et al., 2010
Syzygium cumini	Leaves, seeds	Mycaminose	Ethanollic	Diabetic rats	Kumar et al., 2013

Table 3: A list of recently scrutinized polyherbal formulations.

Formulation with scientific names	Scientific evaluation	Outcome of the study	Reference
Alstonia congensis, Stachytarpheta angustifolia, Xylophia aethiopica	alloxan-induced diabetic rats	Potent hypoglycaemic activity and beneficial effects on cardiovascular risk factors	Ogbonnia et al., 2010
Achillea millefolium, Agathosma betulina, Salvia officinalis, Taraxacum officinalis, Thymus vulgaris, Trigonella foenum-graecum, Urtica urens	In vitro study	strong antioxidant and antidiabetic activity	Paddy et al., 2015
Allium sativum, Cinnamomum zeylanicum, Citrullus colocynthis, Juglans regia, Nigella sativa, Olea europaea, Punica granatum, Salvia officinalis, Teucrium polium, Trigonella foenum, Urtica dioica, Vaccinium arctostaphylos	Streptozotocin induced diabetic rats	Beneficial effects on blood glucose and lipid profile and has a potential to be used as a dietary supplement for the management of diabetes	Ghorbani et al., 2013
Allium sativum, Zingiber officinale, Capsicum frutescens	alloxan-induced diabetic rats	Recommended for the treatment of diabetes	Otunola et al., 2015
Glycosmis pentaphylla, Tridax procumbens, Mangifera indica	Streptozotocin induced diabetic rats	significant antidiabetic activity	Petchi et al., 2014
Eugenia jambolana, Gymnema sylvestre, Momordica charantia and Andrographis paniculata, Myristica fragrans	Streptozotocin induced diabetic rats	Significantly reduced blood glucose level	Nagja et al., 2017

Toxicity of medicinal plants and polyherbal mixtures

Toxicology is a branch of science that deals with toxic substances and their effects. Toxicological studies are very important for the development of new safe drugs and for the extension of the therapeutic potential of existing drugs (Parasuraman, 2011). Although medicinal plants play a significant role in the treatment of diseases, many unsafe and fatal side effects have also been reported (Ekor, 2014). These toxic effects might be due to allergic reactions, effects from impurities and interactions with drugs and other herbs (Hussain, 2011). The Organization for Economic Co-operation and Development (OECD) and the International Conference on Harmonization (ICH) have established the guidelines for toxicity studies of pharmacological substances including medicinal plant extracts, herbal mixtures and for isolated compounds (Kiani *et al.*, 2018). Toxicological studies determines whether the plant extract can be accepted for clinical use or not. Even though many phytochemicals are commonly used for health benefits, some of them have been reported as toxic substances and may be present in food or herbal remedies. Arseculeratne and co-workers (1985) have reported some Sri Lankan medicinal plants which are popular in Ayurvedic medicine exerted significant toxic effects such as *Terminalia chebula*. However, there are many antidiabetic plants extracts which have been documented as toxic free extracts with recent scientific investigations such as *Coccinia grandis* (Attanayake *et al.*, 2013b). Table 4 describes toxicological studies conducted on medicinal plants with

potent antidiabetic activity.

CONCLUSIONS

In the present review we described the traditional antidiabetic herbal medicines with an emphasis on scientific investigations conducted on antidiabetic medicinal plants, polyherbal mixtures, isolated compounds and their associated toxicities. Herbal medicines are used by a large number of patients with diabetes mellitus since ancient times. The indigenous knowledge on herbal medicines can be successfully blended with modern scientific knowledge to develop antidiabetic agents that can be used in the management of diabetes mellitus. The major hindrance in the amalgamation of medicinal plants with modern therapeutic applications is lack of scientific and clinical data proving their efficacy and safety. Many investigations have been carried out to scrutinize the therapeutic benefits of medicinal plants, and polyherbal mixtures used in different traditional medicine systems. However, a considerable number of medicinal plants used in traditional medicines have not been investigated yet. Therefore, there is a crucial need to conduct scientific investigations to scrutinize the therapeutic potency of medicinal plant extracts and polyherbal mixtures in clinical practice, elucidate the antidiabetic principle(s), and to isolate antidiabetic compounds in the near future.

Table 4: Reported toxic effect of plants which are used in traditional medicine.

Plant	Ethanobotanical use	Used plant part	Solvent	Experimental period	Type of toxicity	Reference
<i>Aframomum melegueta</i>	Stomachache, diarrhea	Seed	Ethanol	Sub chronic (28 days)	Liver	Ilic <i>et al.</i> , 2010
<i>Anacardium occidentale</i>	Diabetes mellitus	Leaves	Hexane	Acute (7 days), Chronic (56 days)	Liver and Kidney	Tedong <i>et al.</i> , 2007
<i>Cylicodiscus gabunensis</i>	Gastrointestinal disorders	Stem bark	Ethyl acetate	Acute (7 days), sub-acute (6 weeks)	Liver, Kidney and Lungs	Mabeku <i>et al.</i> , 2007
<i>Drimysan gustifolia</i>	antiulcer and antiinflammatory	Leaves	Ethanol	Acute (48 hours)	Behavioral changes	Gomes <i>et al.</i> , 2013
<i>Galega officinalis</i>	Antidiabetic	Aerial parts	Ethanol	Sub chronic (90 days)	Liver and Lungs	Rasekh <i>et al.</i> , 2008
<i>Mammea africana</i>	Hypercholesterolaemia	Stem bark	Water	Acute (48 hours), sub-acute (21 days)	Liver	Antia <i>et al.</i> , 2006
<i>Poygala fruticos</i>	Chronic ulcer	Leaves	Water	Acute (14 days), Sub chronic (31 days)	anti-haemopoietic, hepatotoxic and nephrotoxic	Mukinda and Eagles, 2010

ACKNOWLEDGEMENT

We would like to acknowledge University Research Grant B/Sc/2016/CH/URG/52/S from the University of Peradeniya, Peradeniya, Sri Lanka for funding the research project.

DECLARATION OF CONFLICT OF INTEREST

The authors declare that no conflicts of interest.

REFERENCES

- Abesundara, K.J., Matsui, T. and Matsumoto, K. (2004). α -Glucosidase inhibitory activity of some Sri Lanka plant extracts, one of which, *Cassia auriculata*, exerts a strong antihyperglycaemic effect in rats comparable to the therapeutic drug acarbose. *Journal of Agricultural and Food Chemistry* **52**: 2541-2545. DOI: <https://doi.org/10.1021/jf035330s>.
- Alabi, A., Ajayi, A.M., Olooto, W.E., Emegoakor, C., Oladunjoye, O. and Obikoya, Y. (2017). Antinociceptive and anti-inflammatory properties of a polyherbal extract of *Plumbago zeylinica* and *Capsicum frutescens* in rodents. *African Journal of Biomedical Research* **20**: 277-285. DOI : <https://www.ajol.info/index.php/ajbr/article/view/163651>.
- Antia, B.S., Okokon, J.E., Nwidu, L.L. and Jackson, C.L. (2006). Effect of subchronic administration of ethanolic stem bark extract of *Mammea africana* Sabine on haematological and biochemical parameters of rats. *African Journal of Biomedical Research* **9**: 129-132. DOI : <https://www.ajol.info/index.php/ajbr/article/view/48894>.
- American Diabetes Association. (2013). Diagnosis and classification of diabetes mellitus, *Diabetes care*, **36**: S67-S74. DOI : <https://doi.org/10.2337/dc13-S067>
- Arambewela, L.S., Arawwawala, L.D. and Ratnasooriya, W.D. (2004). Antinociceptive activities of aqueous and ethanolic extracts of *Alpinia calcarata* rhizomes in rats. *Journal of Ethnopharmacology* **95**: 311-316. DOI : <https://doi.org/10.1016/j.jep.2004.07.015>.
- Arambewela, L.S., Arawwawala, L.D. and Ratnasooriya, W.D. (2005). Antidiabetic activities of aqueous and ethanolic extracts of *Piper betle* leaves in rats. *Journal of Ethnopharmacology* **102**: 239-245. DOI : <https://doi.org/10.1016/j.jep.2005.06.016>.
- Arseculeratne, S.N., Gunatilaka, A.L. and Panabokke, R.G. (1985). Studies on medicinal plants of Sri Lanka. Part 14: toxicity of some traditional medicinal herbs. *Journal of Ethnopharmacology* **13**: 323-333 DOI : [https://doi.org/10.1016/0378-8741\(85\)90078-9](https://doi.org/10.1016/0378-8741(85)90078-9).
- Aswathy, T.S., Jessykutty, P.C. (2017). Antidiabetic phyto resources: A review. *Journal of Medicinal Plants Studies* **5**: 165-169.
- Attanayake, A.P., Jayatilaka, K.A., Pathirana, C. and Mudduwa, L.K. (2013a). Study of antihyperglycaemic activity of medicinal plant extracts in alloxan induced diabetic rats. *Ancient Science of life* **32**: 193-198. DOI : <https://dx.doi.org/10.4103%2F0257-7941.131970>.
- Attanayake, A.P., Jayatilaka, K.A., Pathirana, C. and Mudduwa, L.K. (2013b). Efficacy and toxicological evaluation of *Coccinia grandis* (Cucurbitaceae) extract in male Wistar rats. *Asian Pacific Journal of Tropical Disease* **3**: 460-466. DOI : [https://doi.org/10.1016/S2222-1808\(13\)60101-2](https://doi.org/10.1016/S2222-1808(13)60101-2).
- Attanayake, A.P., Jayatilaka, K.A.P.W., Pathirana, C. and Mudduwa, L.K.B. (2016a). *Gmelina arborea* Roxb. (family: Verbenaceae) extract upregulates the β -cell regeneration in STZ induced diabetic rats. *Journal of Diabetes Research* **2016**: 8. DOI : <https://doi.org/10.1155/2016/4513871>.
- Attanayake, A.P., Jayatilaka, K.A.P.W. and Mudduwa, L.K.B. (2016b). Anti-diabetic potential of ivy gourd (*Coccinia grandis*, family: Cucurbitaceae) grown in Sri Lanka: A review. *J. Pharmacogn Phytochem* **5**: 286-289.
- Bahmani, M., Zargaran, A., Rafieian-Kopaei, M. and Saki, K. (2014). Ethnobotanical study of medicinal plants used in the management of diabetes mellitus in the Urmia, Northwest Iran. *Asian Pacific Journal of Tropical Medicine* **7**: 348-354. DOI : [https://doi.org/10.1016/S1995-7645\(14\)60257-1](https://doi.org/10.1016/S1995-7645(14)60257-1).
- Brahmachari, G. (2011). Bio-flavonoids with promising antidiabetic potentials: A critical survey. *Research Signpost* **661**: 187-212.
- Briskin, D.P. (2000). Medicinal plants and phytomedicines. Linking plant biochemistry and physiology to human health. *Plant Physiology* **124**:507-514. DOI : <https://doi.org/10.1104/pp.124.2.507>.
- Chaudhury, A., Duvoor, C., Dendi, R., Sena, V., Kraleti, S., Chada, A., Ravilla, R., Marco, A., Shekhawat, N.S., Montales, M.T. and Kuriakose, K. (2017). Clinical review of antidiabetic drugs: implications for type 2 diabetes mellitus management. *Frontiers in Endocrinology* **8**: 6. DOI : <https://dx.doi.org/10.3389%2Ffendo.2017.00006>.
- Chang, C.L., Lin, Y., Bartolome, A.P., Chen, Y.C., Chiu, S.C. and Yang, W.C. (2013). Herbal therapies for type 2 diabetes mellitus: chemistry, biology, and potential application of selected plants and compounds. *Evidence-Based Complementary and Alternative Medicine* **2013**: 33. DOI : <https://doi.org/10.1155/2013/378657>.
- Chawla, A., Chawla, R. and Jaggi, S. (2016). Microvascular and macrovascular complications in diabetes mellitus: distinct or continuum?. *Indian Journal of Endocrinology and Metabolism* **20**: 546. DOI : <https://dx.doi.org/10.4103%2F2230-8210.183480>.
- Che, C.T., Wang, Z.J., Chow, M.S. and Lam, C.W. (2013). Herb-herb combination for therapeutic enhancement and advancement: theory, practice and future perspectives. *Molecules* **18**: 5125-5141. DOI : <https://doi.org/10.3390/molecules18055125>.
- Cheng, D., Liang, B. and Li, Y. (2012). Antihyperglycaemic effect of *Ginkgo biloba* extract in streptozotocin-induced diabetes in rats. *BioMed Research International* **2013**: 7.
- Delaviz, H., Mohammadi, J., Ghalamfarsa, G., Mohammadi, B., Farhadi, N. (2017). A review study on phytochemistry and pharmacology applications of *Juglans regia* plant. *Pharmacognosy Reviews* **11**: 145-152. DOI : https://dx.doi.org/10.4103%2Fphrev.phrev_10_17.
- Dinesh kumar, B., Mitra, A. and Mahadevappa, M. (2010).

- Antidiabetic and hypolipidemic effects of mahanimbine (carbazole alkaloid) from *Murraya koenigii* (rutaceae) leaves. *Phytomedicine* **2**: 22-30. DOI:10.5138/ijpm.2010.0975.0185.02004.
- Ediriweera, E.R. and Ratnasooriya, W.D. (2009). A review on herbs used in treatment of diabetes mellitus by Sri Lankan ayurvedic and traditional physicians. *An International Quarterly Journal of Research In Ayurveda* **30**: 373-391.
- Eidi, A., Eidi, M. and Esmaeili, E. (2006). Antidiabetic effect of garlic (*Allium sativum* L.) in normal and streptozotocin-induced diabetic rats. *Phytomedicine* **13**: 624-629. DOI : <https://doi.org/10.1016/j.phymed.2005.09.010>.
- Ekor, M. (2014). The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Frontiers in Pharmacology* **4**: 177. DOI : <https://doi.org/10.3389/fphar.2013.00177>.
- Farzaei, M.H., Rahimi, R., Farzaei, F. and Abdollahi, M. (2015). Traditional medicinal herbs for the management of diabetes and its complications: an evidence-based review. *International Journal of Pharmacology* **11**: 874-87. DOI : <http://dx.doi.org/10.3923/ijp.2015.874.887>.
- Feng, J., Yang, X.W. and Wang, R.F. (2011). Bio-assay guided isolation and identification of α -glucosidase inhibitors from the leaves of *Aquilaria sinensis*. *Phytochemistry* **2**: 242-247. DOI : <https://doi.org/10.1016/j.phytochem.2010.11.025>.
- Fernando, M.R., Wickramasinghe, S.N., Thabrew, M.I., Ariyananda, P.L. and Karunanayake, E.H. (1991). Effect of *Artocarpus heterophyllus* and *Asteracanthus longifolia* on glucose tolerance in normal human subjects and in maturity-onset diabetic patients. *Journal of Ethnopharmacology* **31**: 277-282. DOI: [https://doi.org/10.1016/0378-8741\(91\)90012-3](https://doi.org/10.1016/0378-8741(91)90012-3).
- Ghorbani, A., Shafiee-Nick, R., Rakhshandeh, H. and Borji, A. (2013). Antihyperlipidemic effect of a polyherbal mixture in streptozotocin-induced diabetic rats. *Journal of Lipids* **2013**: 6 DOI : <https://doi.org/10.1155/2013/675759>.
- Gopal, N.M., Tejaswini, J., Mantry, S. and Kumar, S.A. (2014). International standards of medicinal plants. *International Journal of Innovative Pharmaceutical Sciences and Research* **2**: 2498-2532.
- Gothai, S., Ganesan, P., Park, S.Y., Fakurazi, S., Choi, D.K. and Arulselvan, P. (2016). Natural phyto-bioactive compounds for the treatment of type 2 diabetes: inflammation as a target. *Nutrients* **8**:461. DOI : <https://doi.org/10.3390/nu8080461>.
- Grover, J.K., Yadav, S. and Vats, V. (2002). Medicinal plants of India with anti-diabetic potential. *Journal of Ethnopharmacology* **81**: 81-100. DOI : [https://doi.org/10.1016/S0378-8741\(02\)00059-4](https://doi.org/10.1016/S0378-8741(02)00059-4).
- Gunawan-Puteri, M.D. and Kawabata, J. (2010). Novel α -glucosidase inhibitors from *Macaranga tanarius* leaves. *Food Chemistry* **123**: 384-389. DOI : <https://doi.org/10.1016/j.foodchem.2010.04.050>.
- Han, Y., Jung, H.W. and Park, Y.K. (2014). Selective therapeutic effect of *Cornus officinalis* fruits on the damage of different organs in STZ-induced diabetic rats. *The American Journal of Chinese Medicine* **42**: 1169-1182. DOI: <https://doi.org/10.1142/S0192415X14500736>.
- Hannan, J.M., Ali, L., Khaleque, J., Akhter, M., Flatt, P.R. and Abdel-Wahab, Y.H. (2012). Antihyperglycaemic activity of *Asparagus racemosus* roots is partly mediated by inhibition of carbohydrate digestion and absorption, and enhancement of cellular insulin action. *British Journal of Nutrition* **107**: 1316-1323. DOI : <https://doi.org/10.1017/S0007114511004284>.
- Helal, E.G., Abd-Elwahab, S.M., Atia, T.A. and Mohammad, A.A. (2013). Hypoglycaemic effect of the aqueous extracts of *Lupinus albus*, *Medicago sativa* (Seeds) and their mixture on diabetic rats. *Egyptian Journal of Hospital Medicine* **52**: 685-698. DOI : [10.12816/0000607](https://doi.org/10.12816/0000607).
- Hussain, S. (2011). Patient counseling about herbal-drug interactions. *African Journal of Traditional, Complementary and Alternative Medicines* **8**: 152-163. DOI: <https://www.ajol.info/index.php/ajtcam/article/view/67962>.
- Ieyama, T., Gunawan-Puteri, M.D. and Kawabata, J. (2011). α -Glucosidase inhibitors from the bulb of *Eleutherine americana*. *Food Chemistry* **128**: 308-311. DOI: <https://doi.org/10.1016/j.foodchem.2011.03.021>.
- Ilic, N., Schmidt, B.M., Poulev, A. and Raskin, I. (2010). Toxicological evaluation of grains of paradise (*Aframomum melegueta*)[Roscoe] K. Schum. *Journal of Ethnopharmacology* **127**: 352-356. DOI : <https://doi.org/10.1016/j.jep.2009.10.031>.
- Jamshidi-Kia, F., Lorigooini, Z. and Amini-Khoei, H. (2018). Medicinal plants: past history and future perspective. *Journal of Herbmed Pharmacology* **1**: 1-7. DOI : <http://eprints.skums.ac.ir/id/eprint/6978>.
- Jawla, S., Kumar, Y. and Khan, M.S. (2012). Hypoglycaemic activity of *Bougainvillea spectabilis* stem bark in normal and alloxan-induced diabetic rats. *Asian Pacific Journal of Tropical Biomedicine* **2**: 919-23. DOI : [https://doi.org/10.1016/S2221-1691\(12\)60337-2](https://doi.org/10.1016/S2221-1691(12)60337-2).
- Kanaujia, A., Duggar, R., Pannakal, S.T., Yadav, S.S., Katiyar, C.K., Bansal, V., Anand, S., Sujatha, S. and Lakshmi, B.S. (2010). Insulinomimetic activity of two new gallotannins from the fruits of *Capparis moonii*. *Bioorganic & Medicinal Chemistry* **18**: 3940-3945. DOI : <https://doi.org/10.1016/j.bmc.2010.04.032>.
- Karunamoorthi, K., Jegajeevanram, K., Vijayalakshmi, J. and Mengistie, E. (2013). Traditional medicinal plants: a source of phytotherapeutic modality in resource-constrained health care settings. *Journal of Evidence-Based Complementary & Alternative Medicine* **18**: 67-74. DOI : <https://doi.org/10.1177/2156587212460241>.
- Karunanayake, E.H., Welihinda, J., Sirimanne, S.R. and Adorai, G.S. (1984). Oral hypoglycaemic activity of some medicinal plants of Sri Lanka. *Journal of Ethnopharmacology* **11**: 223-231. DOI : [https://doi.org/10.1016/0378-8741\(84\)90040-0](https://doi.org/10.1016/0378-8741(84)90040-0).
- Karunaratne, V., Thadhani, V.M., Khan, S.N. and Choudhary, M.I. (2014). Potent α -glucosidase inhibitors from the lichen *Cladonia* species from Sri Lanka. *Journal of the National Science Foundation of Sri Lanka* **42**: 95-98. DOI : <http://dx.doi.org/10.4038/jnsfsr.v42i1.6684>.
- Kesari, A.N., Gupta, R.K., Singh, S.K., Diwakar, S. and

- Watal, G. (2006). Hypoglycaemic and *Syzygium cumini* antihyperglycaemic activity of *Aegle marmelos* seed extract in normal and diabetic rats. *Journal of Ethnopharmacology* **7**:374-379. DOI : <https://doi.org/10.1016/j.jep.2006.03.042>.
- Kiani, A.A., Ezatpour, B., Niazi, M. and Jahanbakhsh, S. (2018). Toxicity effect of *Nectaroscordum tripedale* extract on hematological and biochemical parameters in mice. *Entomology and Applied Science Letters* **5**:22-25. DOI: <http://eprints.lums.ac.ir/id/eprint/1381>.
- Kim, K.J., Lee, M.S., Jo, K. and Hwang, J.K. (2011). Piperidine alkaloids from *Piper retrofractum* Vahl. protect against high-fat diet-induced obesity by regulating lipid metabolism and activating AMP-activated protein kinase. *Biochemical and Biophysical Research Communications* **411**: 219-225. DOI : <https://doi.org/10.1016/j.bbrc.2011.06.153>.
- Kumar, A., Ilavarasan, R., Deecaraman, M., Aravindan, P., Padmanabhan, N. and Krishan, M.R. (2013). Anti-diabetic activity of *Syzygium cumini* and its isolated compound against streptozotocin-induced diabetic rats. *Journal of Medicinal Plants Research* **2**: 246-249.
- Kumar, P., Baraiya, S., Gaidhani, S.N., Gupta, M.D. and Wanjari, M.M. (2012). Antidiabetic activity of stem bark of *Bauhinia variegata* in alloxan-induced hypoglycaemic rats. *Journal of Pharmacology & Pharmacotherapeutics* **3**: 64-66. DOI: <https://dx.doi.org/10.4103%2F0976-500X.92518>.
- Kumar, R., Pate, D.K., Prasad, S.K., Sairam, K. and Hemalatha, S. (2011). Antidiabetic activity of alcoholic leaves extract of *Alangium lamarckii* Thwaites on streptozotocin–nicotinamide induced type 2 diabetic rats. *Asian Pacific Journal of Tropical Medicine* **4** :904-909. DOI : [https://doi.org/10.1016/S1995-7645\(11\)60216-2](https://doi.org/10.1016/S1995-7645(11)60216-2).
- Kuroda, M., Mimaki, Y., Nishiyama, T., Mae, T., Kishida, H., Tsukagawa, M., Takahashi, K., Kawada, T., Nakagawa, K. and Kitahara, M. (2005). Hypoglycaemic effects of turmeric (*Curcuma longa* L. rhizomes) on genetically diabetic KK-Ay mice. *Biological and Pharmaceutical Bulletin* **28**:937-939. DOI : <https://doi.org/10.1248/bpb.28.937>.
- Mabeku, L.B., Beng, V.P., Kouam, J., Essame, O. and Etoa, F.X. (2007). Toxicological evaluation of ethyl acetate extract of *Cylicodiscus gabunensis* stem bark (Mimosaceae). *Journal of Ethnopharmacology* **111**: 598-606. DOI: <https://doi.org/10.1016/j.jep.2007.01.010>.
- Maroo, J., Vasu, V.T. and Gupta, S. (2003). Dose dependent hypoglycaemic effect of aqueous extract of *Enicostemma littorale* Blume in alloxan induced diabetic rats. *Phytomedicine* **10**: 196-199. DOI: <https://doi.org/10.1078/094471103321659933>.
- Mazid, M., Khan, T.A. and Mohammad, F. (2012). Medicinal plants of rural India: a review of use by Indian folks. *Indo Global Journal of Pharmaceutical Sciences* **2**: 286-304.
- Modak, M., Dixit, P., Londhe, J., Ghaskadbi, S., Devasagayam, T.P. (2007). Recent advances in Indian herbal drug research guest editor: Thomas Paul Asir Devasagayam Indian herbs and herbal drugs used for the treatment of diabetes. *Journal of Clinical Biochemistry and Nutrition* **40**: 163-173. DOI: <https://doi.org/10.3164/jcbrn.40.163>.
- Mukinda, J.T. and Eagles, P.F. (2010). Acute and sub-chronic oral toxicity profiles of the aqueous extract of *Polygala fruticosa* in female mice and rats. *Journal of Ethnopharmacology* **128**: 236-240. DOI: <https://doi.org/10.1016/j.jep.2010.01.022>.
- Muraoka, O., Morikawa, T., Miyake, S., Akaki, J., Ninomiya, K., Pongpiriyadacha, Y. and Yoshikawa, M. (2011). Quantitative analysis of neosalacinol and neokotalanol, another two potent α -glucosidase inhibitors from *Salacia* species, by LC-MS with ion pair chromatography. *Journal of Natural Medicines* **65**:142-148. DOI: <https://doi.org/10.1007/s11418-010-0474-x>.
- Nagja, T., Kumar, V. and Sanjeev, A. (2017). Anti-diabetic activity of a polyherbal formulation in streptozotocin induced type 2 diabetic rats. *Journal of Natural Remedies* **16**: 148-152.
- Ogbonnia, S.O., Mbaka, G.O., Adekunle, A., Anyika, E.N., Gbolade, O.E. and Nwakakwa, N. (2010). Effect of a poly-herbal formulation, Okudiabet, on alloxan-induced diabetic rats. *Agriculture and Biology Journal of North America* **1**:139-145.
- Ogurtsova, K., da Rocha Fernandes, J.D., Huang, Y., Linnenkamp, U., Guariguata, L., Cho, N.H., Cavan, D., Shaw, J.E. and Makaroff, L.E. (2017). IDF Diabetes Atlas: Global estimates for the prevalence of diabetes for 2015 and 2040. *Diabetes Research and Clinical Practice* **128**: 40-50. DOI: <https://doi.org/10.1016/j.diabres.2017.03.024>.
- Utunola, G.A. and Afolayan, A.J. (2015). Antidiabetic effect of combined spices of *Allium sativum*, *Zingiber officinale* and *Capsicum frutescens* in alloxan-induced diabetic rats. *Frontiers in Life Science* **8**: 314-323. DOI: <https://doi.org/10.1080/21553769.2015.1053628>.
- Paddy, V., van Tonder, J.J. and Steenkamp, V. (2015). In vitro antidiabetic activity of a polyherbal tea and its individual ingredients. *Journal of Pharmaceutical Research International* **6**: 389-401. DOI: <https://doi.org/10.9734/BJPR/2015/17583>.
- Parasuraman, S., Thing, G.S. and Dhanaraj, S.A. (2014). Polyherbal formulation: concept of ayurveda. *Pharmacognosy Reviews* **8**: 73-80. DOI: <https://dx.doi.org/10.4103%2F0973-7847.134229>.
- Parasuraman, S. (2011). Toxicological screening. *Journal of Pharmacology & Pharmacotherapeutics* **2**:74. DOI: <https://dx.doi.org/10.4103%2F0976-500X.81895>
- Patel, D.K., Kumar, R., Laloo, D. and Hemalatha, S. (2012). Diabetes mellitus: an overview on its pharmacological aspects and reported medicinal plants having antidiabetic activity. *Asian Pacific Journal of Tropical Biomedicine* **2**: 411-420. DOI: [https://doi.org/10.1016/S2221-1691\(12\)60067-7](https://doi.org/10.1016/S2221-1691(12)60067-7).
- Petchi, R.R., Vijaya, C., and Parasuraman, S. (2014). Antidiabetic activity of polyherbal formulation in streptozotocin–nicotinamide induced diabetic Wistar rats. *Journal of Traditional and Complementary Medicine* **4**: 108-117. DOI: <https://doi.org/10.4103/2225-4110.126174>.

- Rani, M.S., Pippalla, R.S., Mohan, G.K., Gaikwad, S.B. and Prasad, B. (2015). A comparative quality of life assessment of herbal drugs with allopathic drugs for the treatment of diabetic complications. *Pharmaceutical Crops* **5**: 77-84. DOI: <http://dx.doi.org/10.2174/221029060140501077>.
- Rasekh, H.R., Nazari, P., Kamli-Nejad, M. and Hosseinzadeh, L. (2008). Acute and subchronic oral toxicity of *Galega officinalis* in rats. *Journal of Ethnopharmacology*. 116:21-26. DOI: <https://doi.org/10.1016/j.jep.2007.10.030>.
- Rasineni, K., Bellamkonda, R., Singareddy, S.R. and Desireddy, S. (2010). Antihyperglycaemic activity of *Catharanthus roseus* leaf powder in streptozotocin-induced diabetic rats. *Pharmacognosy Research* **2**:195-201. DOI: <https://dx.doi.org/10.4103/2F0974-8490.65523>.
- Sasidharan, S., Chen, Y., Saravanan, D., Sundram, K.M. and Latha, L.Y. (2011). Extraction, isolation and characterization of bioactive compounds from plants extracts. *African Journal of Traditional, Complementary and Alternative Medicines* **8**: 1-10. DOI: <https://www.ajol.info/index.php/ajtcam/article/view/60483>.
- Shailajan, S. (2015). Quality evaluation and standardization of a traditional Unani formulation Jawarish e Amla Sada. *International Journal of Green Pharmacy* **9**: 21-25. DOI: <http://dx.doi.org/10.22377/ijgp.v9i1.432>.
- Shirwaikar, A., Rajendran, K., Kumar, C.D. and Bodla, R. (2004). Antidiabetic activity of aqueous leaf extract of *Annona squamosa* in streptozotocin–nicotinamide type 2 diabetic rats. *Journal of Ethnopharmacology* **91**: 171-175. DOI: <https://doi.org/10.1016/j.jep.2003.12.017>.
- Souza, A., Mbatchi, B. and Herchuelz, A. (2011). Induction of insulin secretion by an aqueous extract of *Tabernanthe iboga* Baill.(Apocynaceae) in rat pancreatic islets of Langerhans. *Journal of Ethnopharmacology* **133**: 1015-1020. DOI: <https://doi.org/10.1016/j.jep.2010.11.028>.
- Sudasinghe, H.P. and Peiris, D.C. (2018). Hypoglycaemic and hypolipidemic activity of aqueous leaf extract of *Passiflora suberosa* L. *Journal of Life and Environmental Sciences* **6**: e4389. DOI: <https://doi.org/10.7717/peerj.4389>.
- Tédong, L., Dzeufiet, P.D., Dimo, T., Asongalem, E.A., Sokeng, S.D., Flejou, J.F., Callard, P. and Kamtchouing, P. (2007). Acute and subchronic toxicity of *Anacardium occidentale* Linn (Anacardiaceae) leaves hexane extract in mice. *African Journal of Traditional, Complementary and Alternative Medicines* **4**: 140-147. DOI: <https://www.ajol.info/index.php/ajtcam/article/view/31194>.
- Thadhani, V.M. and Karunaratne, V. (2017). Potential of lichen compounds as antidiabetic agents with antioxidative properties: A review. *Oxidative Medicine and Cellular Longevity* **2017**: 10. DOI: <https://doi.org/10.1155/2017/2079697>.
- Tian, L.Y., Bai, X., Chen, X.H., Fang, J.B., Liu, S.H. and Chen, J.C. (2010). Anti-diabetic effect of methylswertianin and bellidifolin from *Swertia punicea* Hemsl. and its potential mechanism. *Phytomedicine* **17**:533-539. DOI: <https://doi.org/10.1016/j.phymed.2009.10.007>.
- Trojan-Rodrigues, M., Alves, T.L., Soares, G.L. and Ritter, M.R. (2012). Plants used as antidiabetics in popular medicine in Rio Grande do Sul, southern Brazil. *Journal of Ethnopharmacology* **139**: 155-163. DOI: <https://doi.org/10.1016/j.jep.2011.10.034>.
- Uvarani, C., Jaivel, N., Sankaran, M., Chandraprakash, K., Ata, A. and Mohan, P.S. (2014). Axially chiral bis carbazoles and biological evaluation of the constituents from *Murraya koenigii*. *Fitoterapia* **94**: 10-20. DOI: <https://doi.org/10.1016/j.fitote.2014.01.004>.
- Waisundara, V.Y. and Watawana, M.I. (2014). The classification of Sri Lankan medicinal herbs: An extensive comparison of the antioxidant activities. *Journal of Traditional and Complementary Medicine* **4**: 196-202. DOI: <https://doi.org/10.4103/2225-4110.126175>.
- Wongsa, P., Chaiwarit, J. and Zamaludien, A. (2012). In vitro screening of phenolic compounds, potential inhibition against α -amylase and α -glucosidase of culinary herbs in Thailand. *Food Chemistry* **131**: 964-971. DOI: <https://doi.org/10.1016/j.foodchem.2011.09.088>.
- World Health Organization 2016. *Global report on diabetes*. World Health Organization. DOI: <http://www.ijncd.org/text.asp?2016/1/1/3/184853>.
- Yin, Z., Zhang, W., Feng, F., Zhang, Y. and Kang, W. (2014). α -Glucosidase inhibitors isolated from medicinal plants. *Food Science and Human Wellness* **3**: 136-174. DOI: <https://doi.org/10.1016/j.fshw.2014.11.003>.
- Yuan, H., Ma, Q., Cui, H., Liu, G., Zhao, X., Li, W. and Piao, G. (2017). How can synergism of traditional medicines benefit from network pharmacology?. *Molecules* **22**: 113. DOI: <https://doi.org/10.3390/molecules22071135>.